

Remote Sensing and Modeling: A tool to provide the spatial information for biomass production potential

K. P. Günther, E. Borg, K. Wißkirchen, M. Schroedter-Homscheidt, B. Fichtelmann, J. Gehrung



Deutsches Zentrum
für Luft- und Raumfahrt e.V.
in der Helmholtz-Gemeinschaft




Introduction

Biomass resource assessment today:

1. Regional or continental estimation of biomass (forestry and agriculture) is based on **statistics** including land cover maps
-> **EUROSTAT, FAOSTAT**
2. Use of biomass is forced by the evaluation of **restrictions** e.g. technical and environmental constraints associated with
 - slope (-> digital elevation model)
 - soils (-> map)
 - wildlife habitat (-> map) and other site factors.
3. Currently, there is a lack of **reliable, dynamic data** delineating the location, quantity, and quality of available biomass (-> no differentiation with regard to “**energy plants**”).

Typical biomass potential map



➤ Spatial resolution is limited by the resolution for statistics (in Europe NUTS)



Goal for the future


Biomass resource assessment tomorrow:

1. Use **remote sensing** data to derive high quality information on **land cover / land use** and on **plant status parameter** (leaf area index, temperature,..)
2. Feed a **carbon model** with the remote sensed products to determine the yearly carbon increase (**Net Primary Productivity, NPP**).
3. Find a relation for **biomass ~ NPP**.

Land cover classification (LCC)

Input parameter for carbon modeling

PELCOM 1km pan-European land cover database



PELCOM Land Cover
based on remote sensing data
- AVHRR -

Scale: 1km x 1km
Legend: 16 classes

Reference: 1995

Land cover classification (LCC)

Input parameter for carbon modeling

Umwelt
Bundes
Amt


Für Mensch und Umwelt



Federal Ministry for the
Environment, Nature Conservation
and Nuclear Safety



Deutsches Zentrum
für Luft- und Raumfahrt e.V.



CORINE Land Cover (CLC 2000)

based on remote sensing data

- LANDSAT, SPOT -

Scale: 1 : 100,000

-> 100m x 100m

Legend: 44 classes (Europe)

37 classes (Germany)

Reference year: 2000

ARTIFICIAL SURFACES

URBAN FABRIC

- 111 Continuous urban fabric
- 112 Discontinuous urban fabric

INDUSTRIAL, COMMERCIAL AND TRANSPORT UNITS

- 121 Industrial, commercial and public units
- 122 Road and rail networks and associated land
- 123 Port areas
- 124 Airport

MINES, DUMPS AND CONSTRUCTION SITES

- 131 Mineral extraction sites
- 132 Dump sites
- 133 Construction sites

ARTIFICIAL NON-AGRICULTURAL VEGETATED AREAS

- 141 Green urban areas
- 142 Sport and leisure facilities

AGRICULTURAL AREAS

ARABLE LAND

- 211 Non-irrigated arable land

PERMANENT CROPS

- 221 Vineyards
- 222 Fruit trees and berries plantations

PASTURES

- 231 Pastures

HETEROGENEOUS AGRICULTURAL AREAS

- 242 Complex cultivation patterns
- 243 Land principally occupied by agriculture, with significant areas of natural vegetation

FOREST AND SEMINATURAL AREA

FORESTS

- 311 Broad-leaved forest
- 312 Coniferous forest
- 313 Mixed forest

SCRUBS AND/OR HERBACEOUS VEGETATION

- 321 Natural grassland
- 322 Moors and heathland
- 324 Transitional woodland-scrub

OPEN SPACES WITH LITTLE OR NO VEGETATION

- 331 Beaches, dunes, sand
- 332 Bare rock
- 333 Sparsely vegetated areas
- 334 Burnt areas
- 335 Glaciers and perpetual snow

WETLANDS

INLAND WETLANDS

- 411 Inland marshes
- 412 Peat bogs

COASTAL WETLANDS

- 421 Salt marshes
- 423 Intertidal flats

WATER BODIES

INLAND WATERS

- 511 Water courses
- 512 Water bodies

MARINE WATERS


- 521 Coastal lagoons
- 522 Estuaries
- 523 Sea and ocean



Deutsches Zentrum
für Luft- und Raumfahrt e.V.
in der Helmholtz-Gemeinschaft

Land cover classification (LCC)

Approach for automatic classification



Schematic presentation of the variation of spectral signature of a landscape object for a year, which is additionally disturbed by cloud cover and snow cover.




First conclusion concerning LCC

- In the near future, land cover / **land use** mapping using remote sensing data will be performed in an automatic way by integrating multi-spectral and multi- temporal data sets. Basic tools are under development at DLR.
- Differentiation of agricultural plants is a challenge for the future and will be integrated.
- **Keyword: energy plants for biomass use**

Plant status (Leaf Area Index LAI)

Input parameter for carbon modeling




NDVI:
Normalized Difference
Vegetation Index

LAI:
Leaf Area Index


fPAR:
Fraction of absorbed PAR

Flowchart of data processing for LAI time series

Carbon modeling (BETHY - DLR)




Carbon modeling (BETHY - DLR)



BETHY: Biosphere Energy Transfer Hydrology Model
 Wolfgang Knorr, MPI for Biogeochemistry Jena, 1997
 Klaus Wißkirchen, DLR, 2005



Carbon modeling (BETHY - DLR)




GPP:
Gross Primary Productivity

NPP:
Net Primary Productivity

NPP = GPP – aut. Respiration

Modelled daily cycle of NPP [kgC m⁻² s⁻¹] (red line) at Tharandt

Carbon modeling (BETHY - DLR)




Regional simulation result of
NPP [$\text{gC m}^{-2} \text{y}^{-1}$] for the
region of Tharandt, near
Dresden / Germany.

Spatial resolution is 1.1 km
Time: 1998



Validation of NEP



GPP:
Gross Primary Productivity


NPP:
Net Primary Productivity

NEP:
Netto Ecosystem Productivity

$NPP = GPP - \text{Aut. Respiration}$

$NEP = NPP - \text{Het. Respiration}$


Combining NPP & crop statistics



after Dieter and Englert, 2001

Straw potential

derived from statistics and NPP (as distribution parameter)



From: Julia Gehrung (DLR, University Karlsruhe), Diploma Thesis, 2006





Conclusion

- Remote sensing and carbon modeling delivers maps of Net Primary Productivity (NPP).
- Depending on the spatial resolution of the remote sensing data, local to continental maps of NPP will be available.
- Looking at energy plants for sustainable biomass use, land use maps are necessary indicating the location of energy plants as well as their plant status.